


Accurate Force Fields for Molecular Simulations of Dye Aggregates from First Principles

Maxim Tafipolski, Bernd Engels

Institut für Physikalische und Theoretische Chemie, Universität Würzburg,
Campus Hubland Nord, Emil-Fischer-Str. 42, D-97074 Würzburg, Germany

Accurate intermolecular forces are needed for simulations of aggregates of large organic chromophores, such as perylene bisimide [1], where a highly anisotropic charge distribution due to the extensive π -conjugation calls for a reliable force field since high quality fully ab initio calculations are computationally very demanding [2]. Recent developments in the symmetry adapted perturbation theory (SAPT) [3] allow partitioning of the total intermolecular energy into different physically well-defined contributions (electrostatic, exchange-repulsion, dispersion and induction) against which the corresponding terms in the force field can be separately parametrized. An approach is described to include the missing charge penetration energy term directly into a force field using a sum over pairwise electrostatic energies between spherical atoms as originally suggested by Spackman [4]. This important contribution to the intermolecular potential can be further refined to reproduce the accurate electrostatic energy between monomers in a dimer by allowing for the radial contraction-expansion of atomic charge densities. This new short-range term is supplemented by a long-range electrostatic contribution described with atomic multipoles (up to quadrupoles) based on distributed multipole analysis [5, 6]. The other components of a force field (exchange-repulsion and dispersion) are parametrized to reproduce the accurate data calculated by SAPT(DFT) [7]. As a proof-of-concept, we have derived the force field parameters suitable for modeling intermolecular interactions between polycyclic aromatic hydrocarbons (PAH) [8]. It is shown that it is possible to have a balanced force field suitable for molecular simulations of large molecules avoiding error cancellation to a large extent.



- [1] F. Würthner, *Chem. Commun.*, **2004**, 1564.
- [2] (a) H-M. Zhao, et al. *J. Am. Chem. Soc.*, **2009**, *131*, 15660.
(b) J. Vura-Weis, et al. *J. Am. Chem. Soc.*, **2010**, *132*, 1738.
- [3] K. Szalewicz, K. Patkowski, B. Jeziorski, *Struct. Bond.*, **2005**, *116*, 43.
- [4] M. A. Spackman, *Chem. Phys. Lett.*, **2006**, *418*, 158.
- [5] A. J. Stone, *J. Chem. Theor. Comput.*, **2005**, *1*, 1128.
- [6] J. W. Ponder, et al. *J. Phys. Chem. B*, **2010**, *114*, 2549.
- [7] A. van der Avoird, et al. *Phys. Chem. Chem. Phys.*, **2010**, *12*, 8219.
- [8] M. Tafipolski, B. Engels, *J. Chem. Theory Comput.*, **2011**, *7*, 1791.